

EVITO® T
MAPP 18671

FUNGICIDE

A suspension concentrate formulation containing 250g/l tebuconazole and 180g/l fluoxastrobin

EVITO T is a broad-spectrum systemic fungicide for control of *Sclerotinia sclerotiorum* in oilseed rape

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

EVITO T

A suspension concentrate formulation containing 250 g/l tebuconazole and 180 g/l fluoxastrobin

Warning

**Causes serious eye irritation
Suspected of damaging fertility or the unborn child**

Very toxic to aquatic life with long lasting effects

Obtain special instructions before use
Use personal protective equipment as required

If exposed or concerned: Get medical advice / attention


Collect spillage

Store locked up

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains 1,2-Benzisothiazolin-3-one.
May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.



IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose (litres product/ha)	Maximum number of applications	Latest Timing	Aquatic Buffer Zone
Oilseed rape	0.8	1 per crop	Up to GS71 (10% of pods have reached final size) and 56 days before harvest	5 metres

Other Specific Restrictions

This product must not be applied via hand-held equipment.
Do not apply before 10% of flowers on main raceme open, main raceme elongating (GS61) or 1st May if this occurs after GS61 in the year of harvest.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

Approval Holder and Marketing Company:
ARYSTA LifeScience SAS,
Route d'Artix – BP 80, 64 150 NOGUERES
Tel + 33 (0)5 59 60 92 92.
www.arystalifescience.com

In case of toxic or transport emergency, ring + 44 (0) 1235 239 670 (NCEC) (24 hour).

Batch No. / Production date: See packaging

GB-FLX-017-01 (03/19)

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate. WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice.

IN CASE OF ACCIDENT OR IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

WASH HANDS AND EXPOSED SKIN before meals and after work.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

Environmental Protection

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms, respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, TIGHTLY CLOSED, IN A SAFE PLACE.

KEEP LOCKED UP.

PROTECT FROM FROST

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESISTANCE

The risk management strategy to reduce the risk of resistance developing to EVITO T is based on several measures. Although strategies are presented individually, the integrated use of combinations of different strategies is feasible, beneficial, and often implemented:

- The combination of two active ingredients with different modes of action
- Maintaining the recommended label rate. Do not use reduced doses. It is possible that reducing the dose rate could enhance the development of resistance.

- Maximum of one application per season. Restrict numbers of treatments applied per growing season and apply only when strictly necessary.

- Apply fungicides during good conditions for spraying

- Use as part of spray programs in tank mixtures and sequences with other fungicides with different modes of action

- Integrated disease management. The integrated use of all types of countermeasures against crop disease is not only highly desirable on economic and environmental grounds, but is also a major strategy for avoiding or delaying fungicide resistance. The use of disease resistant crop varieties, biological control agents, and appropriate hygienic practices, such as crop rotation can reduce disease incidence and reduce the need for fungicides applications, thereby decreasing the potential selection of fungicide-resistant pathogens.

For further advice on resistance management contact your agronomist or specialist advisor or visit the FRAG-UK website.

DISEASE CONTROL

EVITO T is recommended for the control of *Sclerotinia sclerotiorum* in Oilseed rape.

CROP SPECIFIC INFORMATION

EVITO T can be safely used on all varieties of oilseed rape.

Timing

Apply as a protectant spray during flowering. The optimum timing is early to mid-flowering (GS 61-65). A second treatment with another authorised product may be required if disease pressure remains high.

Rate of Use

Apply 0.8 L/ha in 200-400 l water/ha. Use the higher volume of water for dense crops.

MIXING AND APPLICATION

Half fill the spray tank with clean water and begin agitation. Add the required quantity of EVITO T and add the remainder of the water.

Continue agitation until spraying is completed.

Wash out sprayer thoroughly after use by rinsing three times with water.

CONDITIONS OF SALE

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling mixing or use or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

**Net Contents :
4x5 L**

7GB1001-0006

1GB0501-0040/C19

