

DIPROSPERO®

MAPP 19188

FUNGICIDE

Contains 500 g/L (45.3 % w/w) Propamocarb and 90 g/L (8.2 % w/w) dimethomorph as a Suspension Concentrate (MAPP 19188)

Systemic fungicide for the control of foliar late blight (*Phytophthora infestans*) on potato.



Toxic to aquatic life with long lasting effects.

Collect spillage.
Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Important Information

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE.

Crops	Max Individual Dose (L product/ha)	Maximum No. of Treatments (per crop)	Latest Time of Application
Potato	2	5	14 days before harvest

Other Specific Restrictions

Cereal crops may be planted as part of a normal rotation following the use of Diprospero. 120 days must elapse after the application of Diprospero before any other crops are planted.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

Manufacturer, Marketing Company and Approval holder:
Arysta LifeScience Benelux sprl
Rue de Renory 26/1, B-4102 Ougrée
Belgium, Tel. +32 4385 9711

24 hour emergency telephone number: +44 (0) 1235 239670

Manufacturing date and Batch No.: see packaging

SAFETY PRECAUTIONS

Operator protection

WASH CONCENTRATE from skin and eyes immediately.
WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.
WHEN USING DO NOT EAT, DRINK OR SMOKE.

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.
WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS OR WARNING

Wait 14 days after last treatment before harvesting the crop.

DISEASES CONTROLLED

Diprospero is a foliar fungicide containing 500 g/L propamocarb hydrochloride and 90 g/L dimethomorph. It can be used for the preventative and curative treatment of foliar late blight (*Phytophthora infestans*) on potato.

CROP-SPECIFIC INFORMATION

RATE OF USE:

The degree of blight pressure should always be considered when planning fungicide applications for blight control.

Diprospero should be used in a disease control programme at 7 day spray intervals and may be applied at 2 L/ha in 200-600 L water per hectare.

Use longer spray intervals when disease pressure is lower. If disease pressure increases (usually later in the season), then the spray interval should be reduced.

CROP VARIETIES:

All varieties of early and maincrop potatoes may be treated with Diprospero. On crops grown for processing, consult your processor before use.

BLIGHT CONTROL PROGRAMMES AND DISEASE RISK

Fungicides for potato blight control are predominantly protective in effect and to achieve best results the disease control programme should begin before blight becomes active in the crop.

Spraying should commence when the crop meets within the row or following the first blight infection period (whichever is the earlier).

Potato blight development and disease risk is generally favoured by warm wet conditions and there are well established systems for identifying high risk periods from crop and meteorological data. Growers should monitor crops and weather conditions to establish risk levels for their crop. Local or regional information on risk periods may also be a guide and is published from a number of sources.

Crops grown under irrigation should also generally be regarded as at high risk and treated at 7 day intervals. Treatment should be applied after irrigation.

Do not apply Diprospero if rainfall is imminent.

APPLICATION

Add the required quantity of Diprospero to a half-filled spray tank with continuous agitation, then add the remaining volume of water. Spray immediately after mixing and continue agitation during spraying.
Immediately after spraying, drain tank completely. Any contamination on the outside of the spraying equipment should be removed by washing with clean water. Rinse inside of tank with clean water and flush through booms and hoses using at least one-third of the spray tank volume. Half fill tank with clean water, agitate and drain tank completely. Repeat this step again.
When disposing of the washings, avoid environmental contamination by spraying on to crop fields away from drainage channels, which feed water courses.

SPRAY QUALITY

Apply Diprospero as a medium spray as defined by the BCPC system.

RESISTANCE

Diprospero contains a mixture of propamocarb (a carbamate fungicide) and dimethomorph (a CAA fungicide).
Up to 5 applications of Diprospero may be made, making up no more than half of the total number of intended late blight sprays per season.
No more than three consecutive applications of a CAA fungicide should be made in any crop.
It should be used in accordance with FRAC and FRAG-UK resistance management advice for CAA mixtures. For further advice on resistance management for potato blight (*Phytophthora infestans*) contact your agronomist or specialist advisor and visit the FRAG-UK and AHDB websites.

CONDITIONS OF SALE

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling mixing or use or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

GB-DPH-012-01 (0620)

NET CONTENTS: **10 L** e

